

When you are done with your homework you should be able to...

- π Determine continuity at a point and continuity on an open interval
- π Determine one-sided limits and continuity on a closed interval
- π Use properties of continuity
- π Understand and use the Intermediate Value Theorem

DEFINITION OF CONTINUITY

CONTINUITY AT A POINT: A function f is continuous at c if the following three conditions are met.

1. $f(c)$ is defined.
2. $\lim_{x \rightarrow c} f(x)$ exists.
3. $\lim_{x \rightarrow c} f(x) = f(c)$.

CONTINUITY ON AN OPEN INTERVAL:

A function is continuous on an open interval (a, b) if it is continuous at each point in the interval. A function that is continuous on the entire real line is everywhere continuous.

1. Draw the graph of the following functions with the given characteristics on the open interval from a to b :
 - a. The function has a removable discontinuity at $x = 0$

 - b. The function has a nonremovable discontinuity at $x = 0$

THEOREM: THE EXISTENCE OF A LIMIT

Let f be a function and let c and L be real numbers. The limit of $f(x)$ as x approaches c is L if and only if

$$\lim_{x \rightarrow c^-} f(x) = L \quad \text{and} \quad \lim_{x \rightarrow c^+} f(x) = L.$$

CONTINUITY ON A CLOSED INTERVAL:


A function f is **continuous on the closed interval $[a, b]$** if it is continuous on the open interval (a, b) and

$$\lim_{x \rightarrow a^+} f(x) = f(a) \quad \text{and} \quad \lim_{x \rightarrow b^-} f(x) = f(b).$$


The function f is **continuous from the right at a** and **continuous from the left at b** .

2. Graph each function and use the definition of continuity to discuss the continuity of each function.

a. $f(x) = \frac{x^2 - 4}{x + 2}$




b. $g(x) = \tan x$



c.

$$y = \begin{cases} |x|, & x \leq 2 \\ -x, & 2 < x < 4 \\ \frac{x^2}{4}, & x \geq 4 \end{cases}$$



THEOREM: PROPERTIES OF CONTINUITY

If b is a real number and f and g are continuous at $x = c$ then the following functions are also continuous at c .

1. Scalar multiple: bf
2. Sum or difference: $f \pm g$
3. Product: fg
4. Quotient: $\frac{f}{g}, g(c) \neq 0$

The following types of functions are continuous at every point in their domains.

1. Polynomial functions: $p(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + x_0$
2. Rational functions: $r(x) = \frac{p(x)}{q(x)}, q(x) \neq 0$
3. Radical functions: $f(x) = \sqrt[n]{x}$
4. Trigonometric functions: $\sin x, \cos x, \tan x, \cot x, \sec x, \csc x$

3. Explain why the following functions are continuous at every point in their domains.

a. $f(x) = \sqrt{x} - \tan x$

Properties used:

b. $f(x) = \frac{5-x}{x \sin x}$

Properties used:

THEOREM: CONTINUITY OF A COMPOSITE FUNCTION

If g is continuous at c and f is continuous at $g(c)$

then the composite function $(f \circ g)(x) = f(g(x))$

is continuous at c .

4. Explain why the following functions are continuous at every point in their domains.

a. $f(x) = \sqrt[3]{x^2 - 8x + 1}$

Properties used:

b. $g(x) = \tan(2x^2)$


Properties used:

THEOREM: THE INTERMEDIATE VALUE THEOREM

If f is continuous on the closed interval $[a, b]$ and k is any number between $f(a)$ and $f(b)$, then there is **at least** one number c in $[a, b]$ such that $f(c) = k$.

5. Consider the function $f(x) = \cos 2x$ on the closed interval $[0, \pi]$.

a. Sketch the graph of f by hand.



b. State the reason why we can apply the mean value theorem (MVT).

c. Use the MVT to find c such that $f(c) = \frac{1}{2}$.